

WET NURSE™

DAIRY KID MILK REPLACER

[22-22-28]

100% MILK PROTEIN

Feeds Act Registration No. 811015

This feed contains added selenium at 0.3 mg/kg

GUARANTEED ANALYSIS

Crude Protein (min.)	22.0%
Crude Protein from milk sources (min.)	22.0%
Crude Fat (min.)	28.0%
Crude Fibre (max.)	0.25%
Calcium (act.)	0.8%
Phosphorus (act.)	0.6%
Sodium (act.)	0.5%
Vitamin A (min.)	40,000 IU/kg
Vitamin D ₃ (min.)	3,000 IU/kg
Vitamin E (min.)	150 IU/kg

INGREDIENTS

A list of ingredients used in this feed may be obtained from the manufacturer or registrant.

NOTE: To maintain freshness, keep partially used bags closed tightly after opening.

SEE REVERSE SIDE FOR MIXING/FEEDING DIRECTIONS

See Bag for Net Weight
See Bag for Lot Number

Manufactured by:



Prairie Micro-Tech Inc.

Head office: (Registrant)
PO Box 26028
2641 Albert Street N
Regina, SK Canada
S4R 8R7

12-550 Trillium Drive
Kitchener, ON
Canada
N2R 1K3

2610A Progressive Way
Abbotsford, BC Canada
V2T 6H9

www.pmtgroup.com

MIXING DIRECTIONS

milk replacer powder	fresh clean potable water	water temperature
150 grams	1000 mls	38° - 45° C

NOTE:

1. Wash and rinse utensils thoroughly after every feeding.
2. When mixing for a 4°C final solution temperature, mix the powder in 38-45°C water then allow to cool to 4°C.

FEEDING DIRECTIONS

Age (Days)	Feedings Per Day	Milk Replacer Solution Feeding Temperature °C	Mixed Milk Replacer Solution Per Feeding MLS	Mixed Milk Replacer Solution Per day MLS
4 - 7	4	38	225	900
8 - 14	3	38	400	1,200
15 - 21	3	38	450	1,350
22 - 28	2	38	750	1,500
29 - 35	2	4 or 38	825	1,650
36 - 42	2	4 or 38	750	1,500

NOTES: Ensure newborn kids get at least 125 mls of colostrum within the first 12 hours of birth. Thereafter, feed 250 - 500 mls/day of appropriate colostrum for 2-3 days after birth. Kids may require assistance in adapting to artificial nipples for 1 to 2 days. Group feeding temperature should be 4°C to limit intake and prevent overeating. Feedings must be evenly spaced. If kids develop scours, maintain the amount of milk replacer solution fed, but reduce the concentration by 25 to 50 % for 1 - 2 days depending on severity of scours. Provide electrolytes, if necessary.

Starting at 1-2 weeks of age, supply kids with free-choice water and palatable creep feed. Offer high quality forage free-choice starting at 3-4 weeks of age. Follow the recommendations of your feed consultant. It is suggested that kids be weaned at not less than 35 days of age or 10-12 kg of body weight or when they are consuming 0.15 kg of solid feed per day.

CAUTION

Directions for use must be carefully followed. Do not feed in association with another feed containing selenium.

LIABILITY DISCLAIMER

Individual results from the use of this product may vary due to management, environmental, genetic, health and sanitation differences. Therefore, Prairie Micro-Tech Inc. does not warrant or guarantee individual results.