

WET NURSE™

PIGLET MILK REPLACER

[24-24-12]

Feeds Act Registration No. 480033

This feed contains added selenium at 0.3 mg/kg

GUARANTEED ANALYSIS

Crude Protein (min.)	24.0%
Crude Protein from milk sources (min.)	24.0%
Crude Fat (min.)	12.0%
Crude Fibre (max.)	0.1%
Calcium (act.)	0.8%
Phosphorus (act.)	0.6%
Sodium (act.)	0.7%
Zinc (act.)	100 mg/kg
Copper (act.)	10 mg/kg
Vitamin A (min.)	20,000 IU/kg
Vitamin D ₃ (min.)	1,500 IU/kg
Vitamin E (min.)	50 IU/kg

INGREDIENTS

A list of ingredients used in this feed may be obtained from the manufacturer or registrant.

NOTE: To maintain freshness, keep partially used bags closed tightly after opening.

SEE REVERSE SIDE FOR MIXING/FEEDING DIRECTIONS

See Bag for Net Weight
See Bag for Lot Number

Manufactured by:



Prairie Micro-Tech Inc.

Head office: (registrant)

PO Box 26028
2641 Albert Street N
Regina, SK Canada
S4R 8R7

12-550 Trillium Drive
Kitchener, ON
Canada
N2R 1K3

2610A Progressive Way
Abbotsford, BC Canada
V2T 6H9

www.pmtgroup.com

MIXING DIRECTIONS

milk replacer powder	fresh clean potable water	water temperature
180 grams	1000 mls	38° - 45° C

NOTES ON MIXING:

- For accurate mixing, use scales to weigh powder
- Always use fresh potable water for mixing milk replacer solution
- Mix only enough solution for one feeding at a time
- Add powder to water (38° - 45°C) using wire whisk to mix thoroughly
- Wash rinse and sanitize utensils thoroughly after every feeding

FEEDING DIRECTIONS

HAND FEEDING BY BOTTLE OR TROUGH

Feed as per the following schedule:

Age days	Milk Replacer Solution mls/feeding	Feeding temperature of solution	Number of feedings per day
0 - 3	Colostrum*	See Notes*	None
0 - 3 (orphans)	60	38°C	12
4 - 7	125	38°C	12
8 - 13*	250	38°C	6
14 - 18*	250	38°C	4
19 - 21*	250	38°C	2
22	Wean		

Notes *

- If possible, leave the piglets with the sow for a minimum of 1 - 3 days after birth to obtain the benefits of colostrum-containing milk.
- When hand feeding by bottle or trough, offer fresh, high quality creep feed and water daily starting at 10 days of age.
- If piglets develop nutritional scours, reduce the amount of milk replacer powder in the solution by 25 - 50% for 1 to 2 days and provide electrolytes. Gradually return to full feeding levels.

CAUTION

Directions for use must be carefully followed.

Consult PMT's website for further information on feeding Wet Nurse™ milk replacers

www.pmtgroup.com

LIABILITY DISCLAIMER

Individual results from the use of this product may vary due to management, environmental, genetic, health and sanitation differences. Therefore, Prairie Micro-Tech Inc. does not warrant or guarantee individual results.